

Theory of Shubnikov-de Haas oscillations around the $\nu = 1/2$ filling factor of the Landau level: Effect of gauge-field fluctuations.

A.G. Aronov^{†,1,2,3}, E. Altshuler², A.D. Mirlin^{1,4} and P. Wölfle¹

¹ *Institut für Theorie der Kondensierten Materie, Universität Karlsruhe, 76128 Karlsruhe, Germany*

² *Department of Condensed Matter Physics, The Weizmann Institute of Science, 76100 Rehovot, Israel*

³ *A.F. Ioffe Physicotechnical Institute, 194021 St.Petersburg, Russia.*

⁴ *Petersburg Nuclear Physics Institute, 188350 Gatchina, St.Petersburg, Russia.*

(December 15, 1994)

We present a theory of magnetooscillations around the $\nu = 1/2$ Landau level filling factor based on a model with a fluctuating Chern–Simons field. The quasiclassical treatment of the problem is appropriate and leads to an unconventional $\exp[-(\pi/\omega_c\tau_{1/2}^*)^4]$ behavior of the amplitude of oscillations. This result is in good qualitative agreement with available experimental data.

PACS numbers:72.15.Lh, 71.25.Hc

Since the discovery of the Fractional Quantum Hall effect (FQHE) [1], the physical properties of a high mobility electron gas subjected to a strong magnetic field are attracting great interest. Laughlin’s theory [2] gives a very good description of the properties of the FQHE states with filling factors $\nu = 1/(2m + 1)$. The subsequently proposed hierarchy scheme [3] explains, in principle, the existence of FQHE states with arbitrary filling factors $\nu = p/q$ (q is odd). However, some drawbacks of this scheme were discovered later. In particular, the FQHE state with $\nu = p/(2p \pm 1)$ appears only on the p -th level of hierarchy, so that the scheme does not explain why these states are experimentally dominating. This discrepancy motivated Jain [4] to propose a different concept based on converting the electrons into composite fermions by attaching to them an even number of flux quanta. Following a similar approach Halperin, Lee and Read [5] developed a theory for the half filled Landau level (see also [6]).

This theory gives an explanation for many experimentally observed properties of the $\nu = 1/2$ state, such as a non-zero value of the longitudinal resistivity, and an anomaly in the surface acoustic wave propagation [7]. It predicts the formation, at half filling, of a metallic state with well defined Fermi surface. This prediction received additional confirmation in recent experiments [8] where a dimensional resonance of the composite fermions was found. From this point of view, the $\nu = p/(2p \pm 1)$ series can be considered as the usual $\nu = p$ Shubnikov–de Haas oscillations (SDHO) for the composite fermions, providing an explanation for the prominence of the above FQHE states. Indeed, the oscillating behavior of the longitudinal resistivity ρ_{xx} near $\nu = 1/2$ is very much reminiscent of its behavior in low magnetic fields where conventional SDHO take place [9,10].

Comparison of the resistivity oscillations near $\nu = 1/2$ and in a weak magnetic field shows, however, not only a similarity in shape, but also an important difference: the amplitude of the SDHO near half filling decreases on approaching $\nu = 1/2$ ($p = \infty$) much faster than it does in weak fields on approaching zero magnetic field $B = 0$ ($\nu = \infty$). As a result, the $\nu = p/(2p \pm 1)$ oscillations vanish at $p = p_{max} = 6 \div 9$ [9,10]. This reflects a difference in the physical properties between the states at $\nu = 1/2$ and $B = 0$, calling thus for a detailed theoretical consideration.

The crucial feature which distinguishes these states is the presence of fluctuations of the fictitious, Chern–Simons (CS), magnetic field [5]. The scattering of the fermions by these fluctuations turns out to be the dominating mechanism of scattering. On the other hand, the problem of a quantum particle in a random magnetic field in 2D was studied in our recent papers [11,12]. We have shown that whereas the transport relaxation time τ_t can be found in the usual way within perturbation theory, the single particle properties of the model are peculiar. In particular we found a Gaussian shape of the broadened Landau levels and unusual expressions for the single particle relaxation time τ_s and for the amplitude of the de Haas–van Alphen oscillations. These results were obtained by using the formalism of path integrals in coordinate space which turns out to be the most appropriate tool in this case. In the present Letter, we apply these methods to the study of magnetooscillations of the conductivity in the FQHE regime near $\nu = 1/2$.

We consider first the zero-temperature limit, when the fluctuations of the fictitious magnetic field are determined by the randomly located impurities [5]. We find the amplitude of the SDHO in this case to be proportional to $\rho_{xx}^{osc} \propto \exp[-(\pi/\omega_c\tau_{1/2}^*)^4]$ where ω_c is the cyclotron frequency and $\tau_{1/2}^*$ plays the role of an effective relaxation time. This result which describes well the experimental data should be contrasted with the usual behavior $\rho_{xx}^{osc} \propto \exp[-\pi/\omega_c\tau]$ that one obtains for the case of the short range random potential.

At finite temperature the particles are scattered additionally by the thermal fluctuations of the magnetic field. We find however that due to the screening in a strong uniform magnetic field, these fluctuations give a contribution to the single particle relaxation rate, which is small compared to usual temperature-dependent factor

$\exp(-2\pi^2 T/\omega_c)$ originating from the Fermi distribution.

We consider a realistic system formed by the 2D electron gas of density n_e and by the positively charged impurities located in a layer separated by a large distance d_s from the electron plane. The statistical transformation attaches to each electron an even number $\tilde{\phi}$ of flux quanta of the CS gauge field. To describe the vicinity of the $\nu = 1/2$ state, we take $\tilde{\phi} = 2$; however the same formalism with $\tilde{\phi} = 4$ can be applied to the $\nu = 1/4$ state.

In the mean field approximation, the statistical magnetic field $B_{1/2} = 4\pi cn_e/e$ cancels exactly the externally applied field B at $\nu = 1/2$. When the filling factor ν is tuned away from $\nu = 1/2$, the effective uniform magnetic field is equal to $B_{eff} = B - B_{1/2}$. For ν close to $1/2$, the number of filled Landau levels of composite fermions $p \gg 1$, so that the problem can be considered quasiclassically. Essential components of the problem, in addition to the uniform magnetic field B_{eff} are the random magnetic field $h(\mathbf{r})$ and random potential $u(\mathbf{r})$. In the quasiclassical approximation, the quantities of interest can be expressed as a sum over classical trajectories. We will treat the random fields in the framework of quasiclassical perturbation theory, neglecting their influence on the classical trajectories. This approximation is valid provided $\omega_c \tau_t \gg 1$. The trajectories are then simply the cyclotron circles in the uniform field B_{eff} .

In a previous paper [12] we used this quasiclassical approach to calculate the de Haas-van Alphen oscillations of the density of states (DOS) in the presence of a random magnetic field in the limit $\omega_c \tau_t \gg 1$. On this condition the conductivity can be written as a sum over periodic orbits in a similar way [13–15]; the resulting expression has the same structure as for the DOS, except for the overall prefactor representing the non-oscillating contribution. We get therefore, in full analogy with [12]

$$\sigma_{xx} = \sigma_{no} \left[1 - 2i \text{Re} \sum_{k=1}^{\infty} \exp \left\{ 2\pi i p k - \frac{1}{2} \langle S_r^2 \rangle k^2 \right\} \right] \quad (1)$$

where σ_{no} denotes the non-oscillating part of σ_{xx} , summation goes over winding numbers k , $p = \frac{1}{2} \frac{e}{c} R_c^2 B_{eff} = 2\pi \frac{cn_e}{e B_{eff}}$, R_c is the cyclotron radius, S_r denotes the contribution to the action induced by random fields along the classical path of winding number $k = 1$, and the angular brackets denote the average over impurity configurations. A good estimate for the amplitude of the oscillations is given by the first harmonic in (1)

$$\sigma_{xx} = \sigma_{no} \left[1 - 2 \cos(2\pi p) \exp \left(-\frac{1}{2} \langle S_r^2 \rangle \right) \right]. \quad (2)$$

At zero temperature, the potential and magnetic fields fluctuations are dominated by the randomly located impurities. Each impurity creates a scalar potential of the form

$$\int (dq) v_0(q) e^{i\mathbf{q}(\mathbf{r}-\mathbf{r}_i)}; \quad v_0(q) = \frac{2\pi e^2}{\epsilon q} e^{-qd_s}, \quad (3)$$

where $\mathbf{r}_i$ is the projection of the impurity position to the 2D plane, ϵ is the dielectric constant, and $(dq) = d^2 q / (2\pi)^2$. This potential gets renormalized due to the screening by fermions and mixing with the CS field. In the random phase approximation (RPA) one gets:

$$A_\mu = (\delta_\mu^\rho - U_{\mu\nu} K^{\nu\rho})^{-1} A_\rho^{(0)}, \quad (4)$$

where we united scalar A_0 and vector $\mathbf{A}$ potentials in a covariant vector A_μ ; the vector $A_\rho^{(0)}$ represents the bare impurity potentials and therefore has only $\rho = 0$ non-zero component. The tensors $U_{\mu\nu}$ and $K^{\nu\rho}$ represent the bare gauge field propagator and the current-density response tensor respectively.

To evaluate eq.(4) we use the Coulomb gauge $\text{div} \mathbf{A} = 0$, go to the momentum space and choose the momentum $\mathbf{q}$ to be directed along the x -axis: $q_x = q$, $q_y = 0$. Then A_μ has only 2 non-zero components corresponding to $\mu = 0, y$, and both K and U become 2×2 matrices [5]:

$$\begin{aligned} K^{\mu\nu}(q) &= \begin{pmatrix} -m^*/2\pi & -iq\sigma_{xy} \\ iq\sigma_{xy} & \chi q^2 - 2i\omega n_e/qk_F \end{pmatrix} \\ U_{\mu\nu}(q) &= \begin{pmatrix} v(q) & 2\pi i\tilde{\phi}/q \\ -2\pi i\tilde{\phi}/q & 0 \end{pmatrix} \\ A_\mu^{(0)}(q) &= \begin{pmatrix} v_0 e^{-i\mathbf{q}\mathbf{r}_i} \\ 0 \end{pmatrix}, \end{aligned} \quad (5)$$

where m^* is the effective mass of fermions, $\chi = 1/12\pi m^*$ is the magnetic susceptibility and $v(q) = 2\pi e^2/(\epsilon q)$ is the Coulomb propagator, and σ_{xy} is the Hall conductivity of composite fermions (not to be confused with electron Hall conductivity).

Substituting (5) in (4), we find

$$A_\mu(q) = \frac{v_0(q) e^{-i\mathbf{q}\mathbf{r}_i}}{\frac{m^* v(q)}{2\pi} + (\tilde{\phi}s + 1)^2 + \frac{\tilde{\phi}^2}{6}} \begin{pmatrix} \tilde{\phi}s + 1 \\ i\tilde{\phi}m^*/q \end{pmatrix}, \quad (6)$$

where $s = 2\pi\sigma_{xy} \simeq p$ in the limit $\omega_c \tau_t \gg 1$.

Let us now compare the first and the second term in denominator of (6). As we will see below, the typical momenta are $q \sim (2d_s)^{-1}$, and we get for $\tilde{\phi} = 2$

$$\frac{m^* v(q)/2\pi}{(2s)^2} = \frac{m^* e^2}{4\epsilon q s^2} \sim \frac{m^* e^2}{\epsilon k_F} \frac{k_F d_s}{2p^2} \sim \frac{50}{p^2}, \quad (7)$$

where $k_F = \sqrt{4\pi n_e}$, and we used typical experimental parameters [9] $n_e = 1.1 \cdot 10^{11} \text{cm}^{-2}$, $d_s = 80 \text{nm}$, and the experimentally estimated value for the ratio $m^* e^2/(\epsilon k_F) \sim 10$. For the not too large p we are interested in, it is thus a reasonable approximation to neglect all but the first term in the denominator of (6). This gives

$$A_\mu(q) = \frac{2\pi}{m^*} \tilde{\phi} e^{-i\mathbf{q}\mathbf{r}_i} e^{-qd_s} \begin{pmatrix} p \\ im^*/q \end{pmatrix}, \quad (8)$$

The random field action S_r in eqs.(1), (2) is given by $S_r = -\oint A_\mu dr^\mu = -(\int A_0 dt - \oint \mathbf{A} d\mathbf{r})$, where the integration goes around a cyclotron orbit. Assuming now the

impurities to be randomly distributed with concentration n_i and uncorrelated, we find

$$\begin{aligned} \langle S_r^2 \rangle &= (2\pi\tilde{\phi})^2 n_i \\ &\times \int (dq) e^{-2qd_s} \left| \frac{p}{k_F} \oint dl e^{-i\mathbf{q}\mathbf{r}} + \int d^2r e^{-i\mathbf{q}\mathbf{r}} \right|^2 \\ &= n_i (4\pi^2 \tilde{\phi} R_c)^2 \int (dq) e^{-2qd_s} \left| \frac{p}{k_F} J_0(qR_c) + \frac{1}{q} J_1(qR_c) \right|^2 \end{aligned} \quad (9)$$

Here $\oint dl$ means integration along the cyclotron orbit and corresponds to the electric field contribution, whereas $\int d^2r$ goes over the area surrounded by the orbit and describes the magnetic field contribution. Taking into account that $R_c^2 = p^2/(\pi n_e)$, we have $R_c/2d_s = p/\sqrt{4\pi n_e d_s^2} \sim p/10 \lesssim 1$. Thus for relevant momenta $q \sim 1/(2d_s)$ and level numbers $p, qR_c \ll 1$ is a reasonable approximation. In this case eq.(9) reduces to

$$\frac{1}{2} \langle S_r^2 \rangle = \pi^3 \tilde{\phi}^2 n_i \frac{R_c^4}{d_s^2} = \frac{n_i}{n_e} \frac{\pi \tilde{\phi}^2}{n_e d_s^2} \left(\frac{2\pi n_e}{m^* \omega_c} \right)^4. \quad (10)$$

Note that electric and magnetic field fluctuations give the same contribution in this limit. According to (2), this gives for the oscillating part of the conductivity:

$$\sigma_{xx}^{osc} \propto -\cos \left(\frac{4\pi^2 n_e c}{e B_{eff}} \right) \exp \left[- \left(\frac{\pi}{\omega_c \tau_{1/2}^*} \right)^4 \right] \quad (11)$$

where we introduced a parameter $\tau_{1/2}^*$ which may be interpreted as an effective relaxation time and is given according to eq.(10) by

$$\tau_{1/2}^* \simeq \frac{m^*}{2n_e} \left(\frac{n_e d_s^2}{4\pi} \right)^{1/4}. \quad (12)$$

When writing eq.(12) we made the usual assumption that concentrations of donors and charge carriers coincide: $n_e = n_i$.

The dependence of the amplitude of oscillations on ω_c in eq.(11) differs from the conventional form $\exp(-\pi/\omega_c \tau)$ which holds for short range potential scattering. We have already shown in [12] that short range magnetic field scattering leads to damping of oscillations in the DOS $\sim \exp[-(\pi/\omega_c \tau)^2]$. As we see now from eq.(11), a long range correlated magnetic field leads to the novel result $\sim \exp[-(\pi/\omega_c \tau)^4]$.

In Fig.1 we present experimental data for the amplitude of ρ^{osc} from [9] ($T = 0.19K$, $B_{eff} > 0$). It is seen that they can be fitted well by $\exp[-(\pi/\omega_c \tau)^4]$, whereas a $\exp(-\pi/\omega_c \tau)$ fit is much worse.

Now we consider the effect of dynamical fluctuations of the gauge field at higher temperatures T . At first glance, one could expect those to be important for the following reason. In the quasistatic approximation, the amplitude of magnetic field fluctuations is given by [17]

$$\langle h(\mathbf{r}) h(\mathbf{r}') \rangle = \delta(\mathbf{r} - \mathbf{r}') T / \chi. \quad (13)$$

This would lead to a suppression of oscillations by a factor $\sim \exp(-\frac{1}{2}\pi R_c^2 T / \chi) \sim \exp(-12\pi^2 p T / \omega_c)$ which could become dominant at high enough temperature. However, eq.(13) was obtained in the absence of a uniform magnetic field, whereas we are considering here the case of a relatively strong field B_{eff} . As we will see below, this leads to a considerable weakening of magnetic field fluctuations.

The propagator of gauge field fluctuations is given by

$$D_{\mu\nu}(q, \omega) = U_{\mu\rho}(q) (\delta_{\rho\nu} - K^{\rho\lambda}(q, \omega) U_{\lambda\nu}(q))^{-1} \quad (14)$$

where $U_{\mu\nu}$ and $K_{\mu\nu}$ were given in eq.(5). In particular for the D_{11} component determining the magnetic field fluctuations, we get

$$\begin{aligned} D_{11}(q, \omega) &= \left(i\omega \frac{2n_e}{qk_F} - \tilde{\chi} q^2 \right)^{-1}; \\ \tilde{\chi} &= \frac{1}{2\pi m^*} \left[\frac{1}{6} + \left(s + \frac{1}{\phi} \right)^2 \right] + \frac{v(q)}{(2\pi\tilde{\phi})^2}. \end{aligned} \quad (15)$$

We see that due to the Hall conductivity $s = 2\pi\sigma_{xy}$, the value of $\tilde{\chi}$ exceeds considerably the bare value of susceptibility $\chi = 1/12\pi m^*$. An analogous suppression of the gauge field fluctuations in the external magnetic field was found by Ioffe and Wiegmann [16] who considered the magnetoresistance of doped Mott insulators.

In the quasistatic approximation we find

$$\langle A_1 A_1 \rangle_q = \int \frac{d\omega}{2\pi} \frac{2T}{\omega} \text{Im} D_{11} = \frac{T}{\tilde{\chi} q^2} = \frac{2\pi m^*}{s^2} \frac{T}{q^2} \quad (16)$$

This leads to the following result for the amplitude of magnetic field fluctuations:

$$\langle hh \rangle_q = \frac{2\pi m^*}{s^2} T \quad (17)$$

The contribution of these fluctuations to the random field action $\langle S_r^2 \rangle$ in (2) is therefore equal to

$$\frac{1}{2} \langle S_r^2 \rangle_T = \frac{1}{2} \pi R_c^2 \frac{2\pi m^*}{s^2} T = \frac{2\pi^2 T}{p\omega_c} \quad (18)$$

Thus the effect of dynamic fluctuations of magnetic field to the amplitude of oscillations is small compared to the usual Fermi factor $(2\pi^2 T / \omega_c) / \sinh(2\pi^2 T / \omega_c)$ and can be neglected.

For completeness, we consider now the SDHO in low fields (around $B = 0$), where there is no CS gauge field. At low temperature the scattering is then due to screened impurity potential [5] $A_0 = (2\pi/m_b) e^{-qd_s} e^{-i\mathbf{q}\mathbf{r}_i}$, where m_b is the band electron mass. (It is easy to see that this follows from eq.(6) if one puts $\tilde{\phi} = 0$.) We find that $\langle S_r^2 \rangle$ in eq.(2) is given for this case by

$$\langle S_r^2 \rangle = (2\pi)^2 \frac{n_i}{n_e} R_c^2 \int q dq e^{-2qd_s} J_0^2(qR_c) \quad (19)$$

Assuming again $n_i = n_e$ and $R_c \ll d_s$, we get for the amplitude of oscillations

$$\sigma_{xx}^{osc} \propto -\cos\left(\frac{2\pi^2 n_e c}{eB}\right) \exp\left[-\left(\frac{\pi}{\omega_c \tau_0^*}\right)^2\right], \quad (20)$$

$$\tau_0^* = \frac{m_b}{n_e} \left(\frac{n_e d_s^2}{\pi}\right)^{1/2}. \quad (21)$$

As was already mentioned, the available experimental data around $\nu = 1/2$ [9] apparently show the behavior $\ln \rho^{osc} \sim 1/\omega_c^4$ predicted by eq.(11). The value of the parameter $\tau_{1/2}^*$ which is found from such a fit (Fig.2) is $\tau_{1/2}^* \simeq 16 \cdot 10^{-12} s$. At the same time the theoretical estimate according to eq.(12) (with use of the parameters of [9]) gives $\tau_{1/2}^* \simeq 2 \cdot 10^{-12} s$ if one uses the experimental value of the effective mass $m^* = 0.7m_e$. A similar discrepancy is found for the low-field relaxation time: eq.(21) for $m_b = 0.07m_e$ gives $\tau_0^* \simeq 1 \cdot 10^{-12} s$, whereas the value quoted in [9] is $\tau_0 \simeq 9 \cdot 10^{-12} s$. Comparison of these values should be taken with certain caution, since τ_0 was found in [9] by using a conventional linear fit for the Dingle plot, rather than eq.(20). We do not expect, however, that this could change τ_0 considerably. We note also that the theoretically estimated values for the transport relaxation rate at $\nu = 1/2$ are typically 4 times greater than extracted from experimental mobilities [5,9]. Therefore the theory seems to overestimate relaxation rates systematically. This situation is not new and has been discussed previously [18,19]. The considerable increase of relaxation times can be attributed to the correlations in positions of charged impurities due to their mutual Coulomb interaction [19]. It is not clear to us, however, whether the here obtained rather sizeable discrepancy between experimental and theoretical values of $\tau_{1/2}^*$ can be explained in this way or else implies an inconsistency of the underlying theoretical picture of $\nu = 1/2$ FQHE.

In conclusion, we have presented a theory of magnetooscillations around $\nu = 1/2$ Landau level filling factor based on a model with fluctuating Chern-Simons field [5]. The quasiclassical treatment of the problem is appropriate and leads to unconventional $\exp\left[-(\pi/\omega_c \tau_{1/2}^*)^4\right]$ behavior of the amplitude of oscillations. This result is in good agreement with available experimental data, although the experimental value of $\tau_{1/2}^*$ exceeds our theoretical estimate.

This work was supported by German-Israel Foundation for Research [GIF, Jerusalem, Israel] (E.A.), WE-Heraeus-Stiftung (A.G.A.), the Alexander von Humboldt Foundation (A.D.M.) and by SFB 195 der Deutschen Forschungsgemeinschaft (P.W.).

[†] Deceased.

- [1] D.C. Tsui, H.L. Stormer and A.C. Gossard, Phys. Rev. Lett. **48**, 1159 (1982).
- [2] R.B. Laughlin, Phys.Rev.Lett. **50**, 1395 (1983).
- [3] F.D.M. Haldane Phys.Rev.Lett. **51**, 605 (1983); B.I. Halperin, Phys.Rev.Lett. **52**, 1583, 2390(E) (1984); R.B. Laughlin, Surface Science, **141**, 11 (1984).
- [4] D.K. Jain, Phys.Rev.Lett. **63**, 199 (1989), Phys.Rev.B **40**, 8079 (1989), Phys.Rev.B **41**, 7653 (1990).
- [5] B.I. Halperin, P.A. Lee, and N. Read, Phys.Rev.B **47**, 7312 (1993).
- [6] V. Kalmeyer and S.-C. Zhang, Phys.Rev.B **46**, 9889 (1992).
- [7] R.L. Willet, M.A. Paalanen, R.R. Ruel, K.W. West, L.N. Pfeiffer, D.J. Bishop, Phys.Rev.Lett. **65**, 112 (1990).
- [8] R.L. Willet, R.R. Ruel, K.W. West, L.N. Pfeiffer, Phys. Rev. Lett. **71**, 3846 (1993); W. Kang, H.L. Stormer, L.N. Pfeiffer, K.W. Baldwin, K.W. West, Phys.Rev.Lett. **71**, 3850 (1993).
- [9] R.R. Du, H.L. Stormer, D.C. Tsui, L.N. Pfeiffer, K.W. West, Solid State Com., **90**, 71 (1994); Phys.Rev.Lett. **70**, 2944 (1993).
- [10] D.R. Leadley, R.J. Nicholas, C.T. Foxon, J.J. Harris, Phys.Rev.Lett. **72**, 1906 (1994).
- [11] A.G. Aronov, A.D. Mirlin and P. Wölfle, Phys.Rev. B **49**, 16609 (1994).
- [12] A.G. Aronov, E. Altshuler, A.D. Mirlin, P.Wölfle, Europhysics Lett., in press.
- [13] A.A. Abrikosov, *Introduction to the theory of normal metals* (Academic Press, New York, 1972).
- [14] G. Hackenbroich and F. von Oppen, to be published.
- [15] K. Richter, to be published.
- [16] L.B. Ioffe, P. Wiegmann, Phys.Rev.B **45**, 519 (1992).
- [17] P.A. Lee and N. Nagaosa, Phys.Rev.B **46**, 5621 (1992).
- [18] P.T. Coleridge, Phys.Rev.B **44**, 3793 (1991).
- [19] E.Buks, M. Heiblum, Y. Levinson, H. Strikman, Semi-cond.Sci.Technol., **9**, 2031 (1994).

FIG. 1. Dingle plot. Logarithm of normalized amplitude of resistivity oscillations $\ln(D_T \rho^{osc}/\rho)$, with $D_T = \sinh(2\pi^2 T/\omega_c)/(2\pi^2 T/\omega_c)$, as a function of inverse effective magnetic field B_{eff}^{-1} . Linear, quadratic and quartic fits of experimental data from [9] are presented.

